



Gender differences in aggression and cortisol levels in zebrafish subjected to unpredictable chronic stress



Cassiano L. Rambo^a, Ricieri Mocelin^b, Matheus Marcon^b, Débora Villanova^b, Gessi Koakoski^c, Murilo S. de Abreu^c, Thiago A. Oliveira^c, Leonardo J.G. Barcellos^{c,d}, Angelo L. Piato^b, Carla D. Bonan^{a,*}

^a PUCRS, Programa de Pós-Graduação em Biologia Celular e Molecular, Laboratório de Neuroquímica e Psicofarmacologia, Av. Ipiranga, 6681, 90619-900 Porto Alegre, RS, Brazil

^b Universidade Federal do Rio Grande do Sul, Programa de Pós-Graduação em Neurociências, Laboratório de Psicofarmacologia e Comportamento, Av. Sarmento Leite 500/107, 90050-170 Porto Alegre, RS, Brazil

^c Universidade Federal de Santa Maria, Programa de Pós-Graduação em Farmacologia, Campus Universitário, Camobi, 97105-900 Santa Maria, RS, Brazil

^d Universidade de Passo Fundo, Programa de Pós-Graduação em Bioexperimentação, BR 285, 99052-900, Passo Fundo, RS, Brazil

HIGHLIGHTS

- UCS increased aggression behavior in males, but not in females.
- Females displayed more aggressive behavior at baseline than control males.
- Stressed females showed increased locomotion compared to stressed and control males.
- Stressed males had higher cortisol levels compared to all other groups.
- No differences in cortisol levels were found between stressed and control females.

ARTICLE INFO

Article history:

Received 24 July 2016

Received in revised form 7 November 2016

Accepted 23 December 2016

Available online 28 December 2016

Keywords:

Aggression

Cortisol

Gender differences

Unpredictable chronic stress

ABSTRACT

Chronic stress may cause physical, behavioral and neuropsychiatric changes, affecting the health condition of an individual. Aggression is a universal behavior with great relevance on human and animal social systems. Despite studies showing the influence of chronic stress on aggression, the effects of unpredictable chronic stress (UCS) on aggressive behavior in male and female zebrafish remain unknown. Thus, the aim of this study was to evaluate the effects of UCS on the aggressive behavior and cortisol levels in adult zebrafish of both sexes. Our results showed that UCS increased aggression in males, but not in females, which displayed more aggressive behavior at baseline than control males. Increased whole-body cortisol levels were observed in stressed males; however, no differences were found between female groups. In conclusion, we reported for the first time gender differences on behavioral parameters and cortisol levels in response to UCS in zebrafish. These results highlight the relevance of studying behavioral and physiological parameters in both sexes separately.

© 2016 Elsevier Inc. All rights reserved.

1. Introduction

Stress is characterized as a process whereby the organism reacts to the threat or stressor by physiological and behavioral changes that, largely, affect the health condition of an individual [5,18]. Stress can induce behavioral and metabolic changes, which may result in somatic and mental disorders [5]. Chronic stress may cause anxiety, depression, executive and/or cognitive dysfunction, atherosclerotic cardiovascular

disease, metabolic disorders, neurovascular degenerative disease, osteopenia, osteoporosis and sleep disorders [7,20]. Thus, excessive stressors and/or chronic exposure can harm a number of essential physiological functions, such as metabolism, growth, reproduction and immune competence, as well as the development of personality and aggressive behavior [17,18,23,30,39].

Aggression is a universal behavior which influences both human and animal social systems [10,11,16,36,37]. This behavior typically occurs in the competition for resources, such as food, mate selection and nesting sites, and is important for protecting offspring and establishing territories and dominance hierarchies [1,10,11,19,24,36–38]. However, intensified aggression can cause harm to an individual and others [16, 25], meaning that this behavior must be carefully controlled [16]. Thus, aggression is a complex set of adaptive behaviors [25,27] that

* Corresponding author at: Laboratório de Neuroquímica e Psicofarmacologia, Programa de Pós-Graduação em Biologia Celular e Molecular, Pontifícia Universidade Católica do Rio Grande do Sul, Av. Ipiranga, 6681, Prédio 12D, sala 301, 90619-900 Porto Alegre, RS, Brazil.

E-mail address: cbonan@pucrs.br (C.D. Bonan).

may have important effects on both the success and evolutionary fitness of individuals [24,28,36].

The zebrafish is a social fish that forms dominance hierarchies in both sexes [32,38]. In this species, aggression is commonly used to monopolize resources like food and mates, occupy nesting sites and protect their status [27,32]. Both males and females may exhibit aggressive behavior, including erection of dorsal, pectoral, caudal and anal fins, dancing or undulating movements, attacks towards the opponent and bites [12,19,38]. They can also display visible color changes after the establishment of dominance, during the fight or during spawning by getting darker and/or more colored [11,12]. This model organism has been used to evaluate the effects of unpredictable chronic stress (UCS) on behavioral, molecular and physiological parameters [33]; however, the effects of UCS on aggressive behavior in male and female zebrafish remain unknown.

Thus, here we evaluate the effects of UCS on aggressive behavior and whole-body cortisol of male and female zebrafish.

2. Materials and Methods

2.1. Animals

A total of 128 adult male ($n = 64$) and female ($n = 64$) 6–8 months old wild-type zebrafish (*Danio rerio*) were obtained from a local commercial supplier (Red Fish, Porto Alegre, Brazil) and acclimated for at least 2 weeks in the experimental room before the experiments. The animals were housed in a 30-L thermostated aquarium filled with unchlorinated water constantly aerated at a targeted temperature of 26 ± 2 °C. Fish were kept under a 14-h light/10-h dark cycle photoperiod and fed twice a day with commercial flake fish food (Alcon BASIC®, Alcon, Brazil). All protocols were approved by the Institutional Animal Care Committee (14/00419, CEUA-PUCRS).

2.2. Unpredictable chronic stress (UCS)

Males and females were divided in two groups: control and stressed. The control and stressed male and female groups were divided in two tanks for each group to minimize a possible tank effect. UCS was performed according to the protocol previously established by Piato et al. [33]. Following a two-week habituation period, the stressed group was subjected twice a day to one of the following stressors during 7 days (Table 1): crowding of 10 animals for 50 min in a 250 mL beaker; social isolation, maintaining animals alone for 45 min in a 250 mL beaker; tank change, three consecutive times; cooling tank water up to 23 °C for 30 min; low water level on housing tanks until animals' dorsal body wall was exposed for 2 min; heating tank water up to 33 °C for 30 min; and chasing animals for 8 min with a net. To prevent habituation and maintain unpredictability, time and sequence of stressors presentation were changed daily. Except during heating and cooling stress, the temperature was controlled during each stressor. Despite the stressful conditions intermittently presented to the fish, no extreme suffering or deaths were observed. A control group (non-stressed) remained in the same room during the equivalent 7-day period.

2.3. Aggression

Twenty-four hours after the UCS, control and stressed fish were subjected to mirror-induced aggression (MIA) protocol previously described by Gerlai [11]. Each fish was individually netted into a small experimental tank ($30 \times 15 \times 10$ cm length \times height \times width). A mirror was placed inclined at 22.5 degrees to the back wall of the tank so that the left vertical edge of the mirror was touching the side of the tank and the right edge was further away. After 30 s for habituation, experimental fish was recorded for 11 min and later analyzed by the ANY-Maze tracking software (Stoelting Co., USA). For the data analysis, the tank was divided in four equal sections and the amount of time spent in the segment closer to the mirror was quantified (Segment 1). In addition, locomotor parameters such as total distance traveled (m), mean speed (m/s) and number of crossings in the tank were evaluated.

2.4. Cortisol extraction and analysis

After aggressive interaction with a mirror, fish were captured and immediately frozen in liquid nitrogen for 10–30 s, followed by storage at -20 °C until cortisol extraction. A pool of 2 fish was used per sample. Whole-body cortisol was extracted using the method previously described by Oliveira et al. [29]. Fish were weighed, minced and placed in a disposable stomacher bag with 2-mL phosphate buffered saline (PBS, pH 7.4) for 6 min. The contents were then transferred to a 10-mL screw top disposable test tube, to which 5 mL of laboratory grade ethyl ether was added. The tube was vortexed for 1 min and centrifuged for 10 min at 3000 rpm, after which the sample was immediately frozen in liquid nitrogen. The unfrozen portion (ethyl ether containing cortisol) was decanted and transferred to a new tube and completely evaporated under a gentle stream of nitrogen for 2 h, yielding a lipid extract containing the cortisol, which was stored at -20 °C. Measurement accuracy was evaluated by calculating the levels recovered from samples spiked with known amounts of cortisol (50, 25 and 12.5 ng/mL). The mean detection of spiked samples was 94.3%. All cortisol values were adjusted for recovery using the following equation: cortisol value = measured value \times 1.0604. Tissue extracts were re-suspended in 1 mL PBS, and whole-body cortisol levels were measured in duplicate samples of each extract using a commercially available enzyme-linked immunosorbent assay kit (EIAgen™CORTISOL test, BioChem ImmunoSystems). The accuracy was tested by repeating the assay 12 times using seven randomly chosen samples on the same plate and calculating the intra-assay coefficient of variation (CV). The reproducibility was tested by assaying the same samples on different plates and calculating the inter-assay CV. To test for linearity and parallelism, the tissue samples were subjected to serial dilutions in the buffer provided with the kit. A strong positive correlation between the curves was observed ($R^2 = 0.8918$), and the samples yielded low inter- and intra-assay CV values (7–10% and 5–9%, respectively).

2.5. Statistical analysis

The data are expressed as mean \pm standard error of mean (S.E.M.). Results were analyzed by two-way ANOVA followed by Tukey's post

Table 1
Procedure of the unpredictable chronic stress protocol in males and females zebrafish.

Monday	Tuesday	Wednesday	Thursday	Friday	Saturday	Sunday
11:50 am	08:00 am	10:30 am	09:15 am	08:45 am	9:30 am	11:30 am
Crowding	Tank change	Low water level	Heating	Social isolation	Cooling	Chasing
4:00 pm	3:30 pm	1:30 pm	5:30 pm	2:30 pm	3:00 pm	4:30 pm
Social isolation	Cooling	Crowding	Chasing	Tank change	Low water level	Heating

hoc test. Two-way ANOVA was used to identify the main effects of gender and UCS, as well as their interactions. Differences were considered significant at $P < 0.05$.

3. Results

Fig. 1 shows the effects of UCS on aggressive behavior and Table 2 summarizes the two-way ANOVA analysis. Stressed males displayed increased aggressive behavior compared to controls, spending more time in the area closer to the mirror (Segment 1). There were no significant differences between control and stressed females in the duration of the time spent in segment 1. However, control females were more aggressive than control males, revealing gender differences in the behavioral response with respect to aggressive behavior.

Fig. 2 shows the effects of gender and UCS on locomotor parameters and Table 2 summarizes the two-way ANOVA analysis. No differences were observed among stressed and controls within same gender in relation to traveled distance (Fig. 2A), mean speed (Fig. 2B), and number of the crossings in tank (Fig. 2C). However, differences between genders were found. Control females showed increased traveled distance (Fig. 2A), mean speed (Fig. 2B), and number of the crossings in tank (Fig. 2C) in relation to control males. Increased traveled distance (Fig. 2A), mean speed (Fig. 2B), and number of the crossings in tank (Fig. 2C) were observed in stressed females compared to stressed and control males.

Fig. 3 shows the effects of gender and UCS on cortisol levels, and Table 2 summarizes the two-way ANOVA analysis. Stressed males showed increased cortisol levels compared to all other groups. There were no differences within the female groups.

4. Discussion

Here we provide evidence that males and females present different stress reactivity and, consequently, differences in aggressive behavior intensity after stress. In fact, females seem to have impaired stress reactivity and, thus, did not respond to an additional acute stressor and did not change their aggressiveness after an acute stress episode. Males seem to be more responsive to an additional acute stressor, probably by a lower response to chronic UCS protocol. Since males typically increase the cortisol levels after this extra acute stressor, they also increase their aggressive behavior in the MIA test.

The basis of our hypothesis of different responses to MIA test after chronic UCS protocol is the feedback mechanism in Hypothalamic-Pituitary-Interrenal (HPI) axis. In fact, female HPI axis seems to be exhausted after the UCS period, incapable to respond to the acute stress provoked by the test. The downregulation of HPI axis after chronic stressors is well reported in fish [3,4], resulting in an impaired cortisol response to subsequent acute stressors. This HPI attenuation might be

Table 2
Results of two-way analysis of variance (ANOVA) for effects of UCS in the aggression behavior.

Dependent variable	Effects	F-value	DF	P-Value
Relative time in Segment 1	UCS	4.192	1, 54	0.0455
	Gender	8.090	1, 54	0.0063
	UCS × Gender	3.877	1, 54	0.0541
Relative time in Segment 1 – analysis across the time	UCS	22.08	3, 594	0.0001
	Gender	1.953	10, 594	0.0361
	UCS × Gender	1.029	30, 594	0.4251
Distance in apparatus	UCS	7.497	1, 54	0.0084
	Gender	32.72	1, 54	0.0001
	UCS × Gender	0.04084	1, 54	0.8406
Mean speed in apparatus	UCS	7.469	1, 54	0.0085
	Gender	32.95	1, 54	0.0001
	UCS × Gender	0.03918	1, 54	0.8438
Line crossings in apparatus	UCS	11.60	1, 54	0.0013
	Gender	22.89	1, 54	0.0001
	UCS × Gender	0.006877	1, 54	0.9342
Whole-body cortisol	UCS	3.098	1, 26	0.0901
	Gender	11.14	1, 26	0.0026
	UCS × Gender	13.89	1, 26	0.0009

a protective mechanism to avoid an over-response in situations of additional acute stressors [34]. The novelty of our study is the clear link between impaired stress reactivity and no changes in aggressive behavior in females after a MIA test.

Aggression is a complex set of behaviors with important effects on the success and evolutionary fitness in human and animal social organizations [16,24,25,27,36] and some studies have shown that chronic stress may influence aggressive behavior [23,30,39].

The link between stress and aggression was proposed due to the correlation between variation in personality traits and in the expression of genes involved in the stress response. Studies have shown that aggression was negatively correlated with cortisol levels and positively correlated with brain expression of glucocorticoid receptors (GR1 and GR2) as a consequence of a physiological pleiotropic effect of glucocorticoid receptors, which are involved in the stress and behavioral responses [2]. Furthermore, the application of an acute stress in zebrafish has a direct correlation to increased aggressive behavior [13].

Differences in aggression patterns were observed in some animal species [40], generally with females more aggressive than males due to its dominant role in social hierarchies [14,22,32,42] and due to mating and social systems [42]. In some fish species, females can be more aggressive, mainly in monogamous or polyandrous systems [21]. However, in zebrafish, differences in aggressive behavior between males and




Fig. 1. Effect of UCS on the aggressive behavior in males and females zebrafish. (A) Relative duration of the time (%) spent in the segment closer of mirror (Segment 1). (B) Relative duration of the time (%) spent in the segment closer of mirror across the time test. The data are presented as the mean \pm S.E.M. Two-way ANOVA followed by Tukey post-hoc test. $n = 14 - 15$. Different letters indicate statistical differences. Statistical significance was set at $P < 0.05$.




Fig. 2. Effect of UCS on the locomotion parameters in males and females zebrafish. (A) Total distance traveled (m) in the apparatus. (B) Mean speed (m/s). (C) Number of crossings. The data are presented as the mean \pm S.E.M. Two-way ANOVA followed by Tukey post-hoc test. $n = 14 - 15$. Different letters indicate statistical differences. Statistical significance was set at $P < 0.05$.

females are not completely clear. Some studies observed equal aggressive behavior in both sexes [1,8,26].

The findings related to cortisol levels and aggression in both male and female control fish reinforce the hypothesis about MIA as an extra stress. Males and females control fish did not present differences in cortisol levels. These cortisol levels in unstressed animals possibly occur due to the 24 h-interval between the end of the UCS protocol and the MIA test, a sufficient time in zebrafish for decreasing the cortisol to the basal levels [35]. On the other hand, females seem to be more aggressive than males, suggesting an effect of UCS protocol. Studies have already observed that females were capable of adapting to repeated social subjugation, even under more severe conditions, whereas males were not able to adapt to these situations [15,40,41]. We consider the MIA test as an additional stressor, since this protocol consists of the transfer fish from the UCS treatment aquarium to the test tank. This transference procedure (persecution, capture, brief air exposure) and the new environment promote physiological changes, such as cortisol increase [31], and anxiety-like behavior [9,13,43].

A possible limitation of our study was the use of four independent groups of fish (males/females vs. UCS/control). This design introduces more variation in the analysis than a repeated measure ANOVA using only two groups (males and females) analyzed before and after the UCS protocol. However, this does not diminish the importance of our data since these findings demonstrated a direct link between gender, stress reactivity and aggression.

It is also important to highlight the relevance and applicability of our results to the general behavioral testing using zebrafish. Gender seems to be an extra source of variation, suggesting the use of sexed populations in studies with behavioral testing. Since psychiatric illness is highly prevalent in modern society [6], it is important to better understand gender differences on physiological responses and activation of HPI-axis in chronic stress and aggression.




Fig. 3. Effect of aggressive behavior on cortisol levels in males and females zebrafish subjected to UCS. The data are presented as the mean \pm S.E.M. Two-way ANOVA followed by Tukey post-hoc test. $n = 7 - 9$. Different letters indicate statistical differences. Statistical significance was set at $P < 0.05$.

Acknowledgements

This work was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (305035/2015-0, 301992/2014-2 and 472715/2012-7).

References

- T.O. Ariyomo, P.J. Watt, Aggression and sex differences in lateralization in the zebrafish, *Anim. Behav.* 86 (2013) 617–622, <http://dx.doi.org/10.1016/j.anbehav.2013.06.019>.
- N. Aubin-Horth, M. Deschênes, S. Cloutier, Natural variation in the molecular stress network correlates with a behavioural syndrome, *Horm. Behav.* 61 (2012) 140–146, <http://dx.doi.org/10.1016/j.yhbeh.2011.11.008>.
- L.J.G. Barcellos, L.C. Kreutz, R.M. Quevedo, Plasmatic levels of cortisol in the response to acute stress in jundiá (*Rhamdia quelen*, Quoy & Gaimard), previously exposed to chronic stress, *Aquaculture* 253 (2006) 317–321.
- L.J.G. Barcellos, S. Nicolayewisky, S.M. Souza, F. Lulhner, Plasmatic levels of cortisol in the response to acute stress in Nile tilapia, *Oreochromis niloticus* (L.), previously exposed to chronic stress, *Aquac. Res.* 30 (1999) 437–444, <http://dx.doi.org/10.1046/j.1365-2109.1999.00348.x>.
- G. Biala, K. Pekala, A. Boguszewska-Czubara, A. Michalak, M. Kruk-Slomka, B. Budzinska, Behavioral and biochemical interaction between nicotine and chronic unpredictable mild stress in mice, *Mol. Neurobiol.* (2016) <http://dx.doi.org/10.1007/s12035-016-9701-0>.
- C.H. Brennan, Zebrafish behavioural assays of translational relevance for the study of psychiatric disease, *Rev. Neurosci.* 22 (2011) 37–48, <http://dx.doi.org/10.1515/RNS.2011.006>.
- G.P. Chrousos, Stress and disorders of the stress system, *Nat. Rev. Endocrinol.* 5 (2009) 374–381, <http://dx.doi.org/10.1038/nrendo.2009.106>.
- S.J. Dahlbom, T. Backström, K. Lundstedt-Enkel, S. Winberg, Aggression and monoamines: effects of sex and social rank in zebrafish (*Danio rerio*), *Behav. Brain Res.* 228 (2012) 333–338, <http://dx.doi.org/10.1016/j.bbr.2011.12.011>.
- R.J. Egan, C.L. Bergner, P.C. Hart, J.M. Cachat, P.R. Canavello, M.F. Elegante, S.I. Elkhayat, B.K. Bartels, A.K. Tien, D.H. Tien, S. Mohnot, E. Beeson, E. Glasgow, H. Amri, Z. Zukowska, A.V. Kalueff, Understanding behavioral and physiological phenotypes of stress and anxiety in zebrafish, *Behav. Brain Res.* 205 (2009) 38–44, <http://dx.doi.org/10.1016/j.bbr.2009.06.022>.
- A.L. Filby, G.C. Paull, T.F. Hickmore, C.R. Tyler, Unravelling the neurophysiological basis of aggression in a fish model, *BMC Genomics* 11 (2010) 498, <http://dx.doi.org/10.1186/1471-2164-11-498>.
- R. Gerlai, Zebra fish: an uncharted behavior genetic model, *Behav. Genet.* 33 (2003) 461–468.
- R. Gerlai, M. Lahav, S. Guo, A. Rosenthal, Drinks like a fish: zebra fish (*Danio rerio*) as a behavior genetic model to study alcohol effects, *Pharmacol. Biochem. Behav.* 67 (2000) 773–782.
- A.C.V.V. Giacomini, M.S. Abreu, L.V. Giacomini, A.M. Siebel, F.F. Zimmerman, C.L. Rambo, R. Mocelin, C.D. Bonan, A.L. Piato, L.J.G. Barcellos, Fluoxetine and diazepam acutely modulate stress induced-behavior, *Behav. Brain Res.* 296 (2016) 301–310, <http://dx.doi.org/10.1016/j.bbr.2015.09.027>.
- L. Goldman, H.H. Swanson, Developmental changes in pre-adult behavior in confined colonies of golden hamsters, *Dev. Psychobiol.* 8 (1975) 137–150, <http://dx.doi.org/10.1002/dev.420080206>.
- K.L. Huhman, M.B. Solomon, M. Janicki, A.C. Harmon, S.M. Lin, J.E. Israel, A.M. Jasnow, Conditioned defeat in male and female Syrian hamsters, *Horm. Behav.* 44 (2003) 293–299.
- L.J. Jones, W.H.J. Norton, Using zebrafish to uncover the genetic and neural basis of aggression, a frequent comorbid symptom of psychiatric disorders, *Behav. Brain Res.* 276 (2015) 171–180, <http://dx.doi.org/10.1016/j.bbr.2014.05.055>.
- R. Kvetnansky, E.L. Sabban, M. Palkovits, Catecholaminergic systems in stress: structural and molecular genetic approaches, *Physiol. Rev.* 89 (2009) 535–606, <http://dx.doi.org/10.1152/physrev.00042.2006>.
- I. Kyrou, C. Tsigos, Stress hormones: physiological stress and regulation of metabolism, *Curr. Opin. Pharmacol.* 9 (2009) 787–793, <http://dx.doi.org/10.1016/j.coph.2009.08.007>.

- [19] E.T. Larson, D.M. O'Malley, R.H. Melloni, Aggression and vasotocin are associated with dominant-subordinate relationships in zebrafish, *Behav. Brain Res.* 167 (2006) 94–102, <http://dx.doi.org/10.1016/j.bbr.2005.08.020>.
- [20] S.J. Lupien, B.S. McEwen, M.R. Gunnar, C. Heim, Effects of stress throughout the lifespan on the brain, behaviour and cognition, *Nat. Rev. Neurosci.* 10 (2009) 434–445, <http://dx.doi.org/10.1038/nrn2639>.
- [21] A.E. Magurran, C.M. Garcia, Sex differences in behaviour as an indirect consequence of mating system, *J. Fish Biol.* 57 (2000) 839–857, <http://dx.doi.org/10.1111/j.1095-8649.2000.tb02196.x>.
- [22] D.M. Marques, E.S. Valenstein, Individual differences in aggressiveness of female hamsters: response to intact and castrated males and to females, *Anim. Behav.* 25 (1977) 131–139.
- [23] B.S. McEwen, Physiology and neurobiology of stress and adaptation: central role of the brain, *Physiol. Rev.* 87 (2007) 873–904, <http://dx.doi.org/10.1152/physrev.00041.2006>.
- [24] K.A. Miczek, E.W. Fish, J.F. De Bold, Neurosteroids, GABA_A receptors, and escalated aggressive behavior, *Horm. Behav.* 44 (2003) 242–257.
- [25] Y.S. Mineur, D.J. Prasol, C. Belzung, W.E. Crusio, Agonistic behavior and unpredictable chronic mild stress in mice, *Behav. Genet.* 33 (2003) 513–519.
- [26] J.A. Moretz, E.P. Martins, B.D. Robison, Behavioral syndromes and the evolution of correlated behavior in zebrafish, *Behav. Ecol.* 18 (2007) 556–562, <http://dx.doi.org/10.1093/beheco/arm011>.
- [27] W. Norton, L. Bally-Cuif, Adult zebrafish as a model organism for behavioural genetics, *BMC Neurosci.* 11 (2010) 90, <http://dx.doi.org/10.1186/1471-2202-11-90>.
- [28] R.F. Oliveira, J.F. Silva, J.M. Simões, Fighting zebrafish: characterization of aggressive behavior and winner-loser effects, *Zebrafish* 8 (2011) 73–81, <http://dx.doi.org/10.1089/zeb.2011.0690>.
- [29] T.A. Oliveira, G. Koakoski, L.C. Kreutz, D. Ferreira, J.G.S. da Rosa, M.S. de Abreu, A.C.V. Giacomini, R.P. Oliveira, M. Fagundes, A.L. Piato, R.E. Barreto, L.J.G. Barcellos, Alcohol impairs predation risk response and communication in zebrafish, *PLoS One* 8 (2013), e75780, <http://dx.doi.org/10.1371/journal.pone.0075780>.
- [30] Ø. Overli, W.J. Korzan, E.T. Larson, S. Winberg, O. Lepage, T.G. Pottinger, K.J. Renner, C.H. Summers, Behavioral and neuroendocrine correlates of displaced aggression in trout, *Horm. Behav.* 45 (2004) 324–329, <http://dx.doi.org/10.1016/j.yhbeh.2004.01.001>.
- [31] N. Pagnussat, A.L. Piato, I.C. Schaefer, M. Blank, A.R. Tamborski, L.D. Guerim, C.D. Bonan, M.R.M. Vianna, D.R. Lara, One for all and all for one: the importance of shoaling on behavioral and stress responses in zebrafish, *Zebrafish* 10 (2013) 338–342, <http://dx.doi.org/10.1089/zeb.2013.0867>.
- [32] G.C. Paull, A.L. Filby, H.G. Giddins, T.S. Coe, P.B. Hamilton, C.R. Tyler, Dominance hierarchies in zebrafish (*Danio rerio*) and their relationship with reproductive success, *Zebrafish* 7 (2010) 109–117, <http://dx.doi.org/10.1089/zeb.2009.0618>.
- [33] Á.L. Piato, K.M. Capiotti, A.R. Tamborski, J.P. Oses, L.J.G. Barcellos, M.R. Bogo, D.R. Lara, M.R. Vianna, C.D. Bonan, Unpredictable chronic stress model in zebrafish (*Danio rerio*): behavioral and physiological responses, *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* 35 (2011) 561–567, <http://dx.doi.org/10.1016/j.pnpbp.2010.12.018>.
- [34] T.G. Pottinger, The effect of stress and exogenous cortisol on receptor-like binding of cortisol in the liver of rainbow trout, *Oncorhynchus mykiss*, *Gen. Comp. Endocrinol.* 78 (1990) 194–203.
- [35] J.M. Ramsay, G.W. Feist, Z.M. Varga, M. Westerfield, M.L. Kent, C.B. Schreck, Whole-body cortisol response of zebrafish to acute net handling stress, *Aquac. Amst. Neth.* 297 (2009) 157–162.
- [36] R.M. Sapolsky, The influence of social hierarchy on primate health, *Science* 308 (2005) 648–652, <http://dx.doi.org/10.1126/science.1106477>.
- [37] J.J. Schwartz, L.A. Ricci, R.H. Melloni, Prior fighting experience increases aggression in Syrian hamsters: implications for a role of dopamine in the winner effect, *Aggress. Behav.* 39 (2013) 290–300, <http://dx.doi.org/10.1002/ab.21476>.
- [38] R. Spence, G. Gerlach, C. Lawrence, C. Smith, The behaviour and ecology of the zebrafish, *Danio rerio*, *Biol. Rev. Camb. Philos. Soc.* 83 (2008) 13–34, <http://dx.doi.org/10.1111/j.1469-185X.2007.00030.x>.
- [39] C.H. Summers, S. Winberg, Interactions between the neural regulation of stress and aggression, *J. Exp. Biol.* 209 (2006) 4581–4589, <http://dx.doi.org/10.1242/jeb.02565>.
- [40] K. Taravosh-Lahn, Y. Delville, Aggressive behavior in female golden hamsters: development and the effect of repeated social stress, *Horm. Behav.* 46 (2004) 428–435, <http://dx.doi.org/10.1016/j.yhbeh.2004.03.007>.
- [41] J.C. Wommack, Y. Delville, Repeated social stress and the development of agonistic behavior: individual differences in coping responses in male golden hamsters, *Physiol. Behav.* 80 (2003) 303–308.
- [42] S.K. Woodley, M.C. Moore, Ovarian hormones influence territorial aggression in free-living female mountain spiny lizards, *Horm. Behav.* 35 (1999) 205–214, <http://dx.doi.org/10.1006/hbeh.1999.1514>.
- [43] L. Ziv, A. Muto, P.J. Schoonheim, S.H. Meijnsing, D. Strasser, H.A. Ingraham, M.J.M. Schaaf, K.R. Yamamoto, H. Baier, An affective disorder in zebrafish with mutation of the glucocorticoid receptor, *Mol. Psychiatry* 18 (2013) 681–691, <http://dx.doi.org/10.1038/mp.2012.64>.