


Sex-dependent metabolic effects of pregestational exercise on prenatally stressed mice

Carolina Luft^{1,2}, Isadora Perez Levides¹, Leonardo Pedrazza³,
Jarbas Rodrigues de Oliveira² and Márcio Vinícius Fagundes Donadio^{1,2} 

Original Article

Cite this article: Luft C, Levides IP, Pedrazza L, de Oliveira JR, and Donadio MVF. (2021) Sex-dependent metabolic effects of pregestational exercise on prenatally stressed mice. *Journal of Developmental Origins of Health and Disease* **12**: 271–279. doi: [10.1017/S2040174420000343](https://doi.org/10.1017/S2040174420000343)

Received: 25 June 2019
Revised: 12 March 2020
Accepted: 8 April 2020
First published online: 14 May 2020

Keywords:

Early life stress; liver; treadmill; sex-difference

Address for correspondence: Professor Márcio V. F. Donadio, Centro Infant, Pontifícia Universidade Católica do Rio Grande do Sul – Av. Ipiranga, 6690, 2º andar, Porto Alegre, Rio Grande do Sul CEP 90610-000, Brazil.
Email: mdonadio@pucrs.br

¹Laboratory of Pediatric Physical Activity, Infant Center, Pontifícia Universidade Católica do Rio Grande do Sul (PUCRS), Porto Alegre, Brazil; ²Laboratory of Cellular Biophysics and Inflammation, Pontifícia Universidade Católica do Rio Grande do Sul (PUCRS), Porto Alegre, Rio Grande do Sul CEP 90619-900, Brazil and ³Laboratory of Ubiquitination and Cellular Signalization, IDIBELL, Campus de Bellvitge, Universitat de Barcelona, L'Hospitalet de Llobregat, ES-08907 Barcelona, Spain

Abstract

Stressful events during the prenatal period have been related to hyperactive hypothalamic–pituitary–adrenal (HPA) axis responses as well as metabolic changes in adult life. Moreover, regular exercise may contribute to the improvement of the symptoms associated with stress and stress-related chronic diseases. Therefore, this study aims to investigate the effects of exercise, before the gestation period, on the metabolic changes induced by prenatal stress in adult mice. Female Balb/c mice were divided into three groups: control (CON), prenatal restraint stress (PNS) and exercise before the gestational period plus PNS (EX + PNS). When adults, the plasmatic biochemical analysis, oxidative stress, gene expression of metabolic-related receptors and sex differences were assessed in the offspring. Prenatal stress decreased neonatal and adult body weight when compared to the pregestational exercise group. Moreover, prenatal stress was associated with reduced body weight in adult males. PNS and EX + PNS females showed decreased hepatic catalase. Pregestational exercise prevented the stress-induced cholesterol increase in females but did not prevent the liver mRNA expression reduction on the peroxisome proliferator-activated receptors (PPARs) α and γ in PNS females. Conversely, PNS and EX + PNS males showed an increased PPAR α mRNA expression. In conclusion, pregestational exercise prevented some effects of prenatal stress on metabolic markers in a sex-specific manner.

Introduction

Maternal health during gestation is an important factor for fetal development. Around 6% of pregnant women report high levels of psychological stress during pregnancy, which results from several conditions such as depression, panic or domestic violence.¹ Higher levels of maternal cortisol may cross over the placental barrier promoting long-lasting effects in the regulation of the hypothalamic–pituitary–adrenal (HPA) axis of the offspring, reducing the expression of glucocorticoid receptors (GR) in the brain and exacerbating glucocorticoid secretion.² Thus, stressful events in the prenatal period have been related to hyperactive HPA axis responses, metabolic changes and low birth weight in the offspring in both early and adult life.^{2–4}

The systemic secretion of adrenal glucocorticoids after stress is central for the mobilization of lipids and glucose reserves, increasing the energy demand to help maintain homeostasis.⁵ For example, individuals with Cushing syndrome or patients receiving treatment with exogenous glucocorticoid administration present hyperglycemia, insulin resistance and increased adiposity.⁶ In the liver, the control of triglycerides and gluconeogenesis is mediated by GR.⁵ In addition, glucocorticoids act to control the regulation of phosphoenolpyruvate carboxykinase (PEPCK), an essential enzyme in the hepatic production of glucose.⁶ Peroxisome proliferator-activated receptors (PPARs) α and γ also play a key role in lipid and glucose metabolism, in addition to oxidation and the storage of fatty acids.⁷ Previous studies on rodents have already demonstrated that prenatal stress is associated with long-term alterations in the offspring's metabolism, with a decreased expression of PPAR α and γ in males and an increased expression of GR in prepubertal mice.^{8,9} These changes may be important in lipid metabolism and glucose-insulin control.

It is already known that regular physical activity is an important non-pharmacological tool to prevent the onset of metabolic disorders^{10,11} and, when performed during pregnancy, may provide benefits to both mother and offspring, both in rodents and humans.¹² We have recently shown that pregestational exercise attenuates the effects of prenatal stress in the offspring, in

Table 1. Cohort characteristics

	Initial number of dams [#]	Final number of dams [#]	Initial number of pups [*]	Final number of pups (adjusted) [*]	Final number of males [*]	Final number of females [*]	Sex ratio
CON	11	8	6.62 ± 1.6	5.37 ± 0.7	2.37 ± 0.9	3.00 ± 0.9	0.79
PNS	10	7	6.85 ± 1.6	5.28 ± 1.5	2.86 ± 1.7	2.43 ± 0.9	1.18
EX + PNS	12	8	5.75 ± 1.6	5.12 ± 0.8	2.62 ± 1.4	2.50 ± 1.2	1.05

Data are shown as absolute number[#] or mean ± standard deviation (SD)^{*}, as indicated. CON, control; PNS, prenatal stress from the second week of pregnancy; EX + PNS, physical exercise three weeks before gestation and prenatal stress from the second week of pregnancy.

a sex-dependent manner.¹³ However, the long-term effects of exercise before pregnancy and the possible mechanisms involved in the metabolic response are not yet well elucidated. Nevertheless, persistent effects of exercise have been related to cognitive improvement in rats¹⁴ and the attenuation of posttraumatic stress symptoms in humans.¹⁵ Acute exercise activates the HPA axis and promotes increased gluconeogenic activity.¹⁶ Physical exercise has also been associated with increased insulin sensitivity in humans with abnormal glucose levels,¹⁷ decreased progression of type 2 diabetes in adults¹⁰ and ameliorated hepatic steatosis in mice.¹¹ The relationship between exercise and HPA axis activity involves a series of complex mechanisms, considering that it may be influenced by duration, type and intensity of exercise, as well as by the characteristics of the stressor and the population studied.

The regulation of stress pathways is markedly influenced by sex differences.¹⁸ Studies with rats have shown that females which are subjected to prenatal stress show greater activation of the HPA axis compared to males.¹⁹ Across the estrous cycle, female mice present different hormonal levels of progesterone and estrogen, which could act directly in the regulation of the physiology and function of several organs and systems.²⁰ Differences between males and females can also be found in metabolic functions and diseases in humans. In general, men have impaired fasting glucose and a greater predisposition to diabetes, while women appear to have a higher incidence of obesity and the accumulation of visceral adiposity.²¹

Considering that fetal exposure to stressful events promote permanent changes in the function of multiple systems and the practice of regular exercise is an important component in the treatment of several diseases, we hypothesized that exercise before pregnancy would contribute to generating beneficial and persistent metabolic effects on the prenatally stressed offspring. Working towards this goal, we investigated the effects of maternal physical exercise before the gestational period on the metabolic changes induced by prenatal stress in adult mice. Body weight, biochemical analyzes, oxidative stress, gene expression of metabolic-related receptors and possible sex differences were evaluated.

Materials and methods

Animals

Male and female Balb/c mice were used to generate the offspring. Animals were kept in individually ventilated boxes, with a light/dark cycle of 12 h (lights on at 07:00 am), at a temperature of 22 ± 2 °C and with free access to water and food. The animals were purchased from the Experimental Biological Models Center (CEMBE) at PUCRS. The experimental protocol was approved by the University Ethics Committee for the Use of Animals (CEUA) under protocol number 15/00446 and all mice were

manipulated according to the guidelines of the National Council for the Control of Animal Experimentation (CONCEA).

Experimental design

Virgin female Balb/c mice (6–8 weeks) were divided into three experimental groups: control (CON, *n* = 8 dams), prenatal stress from the second week of pregnancy (PNS, *n* = 7 dams) and physical exercise 3 weeks before gestation plus prenatal stress from the second week of pregnancy (EX + PNS, *n* = 8 dams). At first, the females had their estrous cycle monitored through a vaginal smear visualization and, during the fertile period, they were mated with males of the same age and lineage. After confirming the mating (presence of a semen plug), females were separated and placed in individual boxes. Four dams were excluded from the EX + PNS group as they presented difficulties in following the exercise protocol (initial number of animals in the group *n* = 12; final number of animals in the group *n* = 8 dams). The cohort characteristics are presented in Table 1. Pregnant females were weighed on days 1, 8 and 15 of the gestation period. The day after birth was considered as postnatal day 1 (PND 1). The litters were randomly adjusted to 4–6 animals per dam on PND 1 and weaned on PND 21. After weaning, males and females were allocated to separate boxes with a maximum of five animals and were maintained until adulthood without any intervention. The offspring were weighed on PND 1, 10 and 60. At 60 days of age (PND 60), female and male mice were euthanized through decapitation, as the use of anesthetics may trigger acute stress responses by activating the HPA axis. The blood was collected for the measurement of plasma glucose, triglycerides and cholesterol concentrations. Moreover, the liver was removed by laparotomy for the evaluation of oxidative stress parameters through the measurement of thiobarbituric acid reactive substances (TBARS), catalase (CAT), reduced glutathione (GSH) and superoxide dismutase (SOD). In addition, glucocorticoid receptor (GR), peroxisome proliferator-activated receptor alpha (PPARα) and peroxisome proliferator-activated receptor gamma (PPARγ) gene expression were evaluated in the liver. The experimental setup is shown in Fig. 1.

Physical exercise

Females of the EX + PNS group were submitted to a protocol of physical exercise on a motorized treadmill, in the last hour of the light cycle, 5 days a week, during the 3 weeks (15 sessions) that preceded the day of mating. The exercise protocol consisted of 60 min of running at a speed of 10 m/min with a 0% slope.²² The exercised animals were adapted to the treadmill for 10 min at a speed of 5 m/min three days before the beginning of the exercise protocol. No stressful stimulus was used to motivate the animals to run. After the end of the protocol, animals were returned to their cages with *ad libitum* access to water and food.




Fig. 1. (a) Experimental design of the study and (b) exercise setup. PNS: prenatal stress from the second week of pregnancy; G-21: 21 days before gestational day 0; G0: gestational day 0; G8: gestational day 8; G21: gestational day 21; PND60: post-natal day 60.

Prenatal stress model

Prenatal restraint stress was applied using the protocol described by Vargas *et al.* (2016).²³ In brief, the protocol was applied from the eighth day of gestation until the day of birth of the offspring, in a closed cylinder (Insight, Brazil), between 10 am and 12 pm, for 30 min, every other day. PNS females were submitted to 5–6 stressful episodes during pregnancy. This prenatal stress model has been previously proven to be a sufficient stressor to increase corticosterone levels in rodents.²⁴ Pregnant control mice remained in their home cages without any intervention during the gestational period.

Biochemical analysis

Biochemical analyses were performed on PND 60 on whole blood samples collected in tubes with heparin and centrifuged (1000g) for 10 min. The separated plasma was then frozen at -80°C until the analyses. The plasma levels of glucose, triglyceride, and cholesterol were determined using commercial kits (Labtest Diagnóstica, Brazil), according to the manufacturer's instructions, in a semi-automatic spectrophotometer (Genesis 8/Spectronic).

Oxidative stress

In order to evaluate the oxidative stress parameters, the liver tissue was homogenized in 10 volumes (1:10, w/v) of sodium phosphate buffer, pH 7.4. The homogenate was centrifuged at 750 g for 10 min at 4°C . The pellet was discarded, and the supernatant was immediately separated and used for the measurements of TBARS, CAT, GSH and SOD. Total protein was measured using the Nanodrop Spectrophotometer (Model 1000, Thermo Scientific).

Thiobarbituric acid reactive substance (TBARS)

The samples were centrifuged, 10 μl of the supernatant was collected and added to 10 μl of sodium dodecyl sulfate (SDS 12.4 mM) and 400 μl of thiobarbituric acid. The mixture was heated for 30 min and then centrifuged at 750 g for 10 min at 25°C . A pink compound was formed and measured spectrophotometrically at 532 nm (Genesis 8/Spectronic). The total content of protein in the sample was measured using the Nanodrop Spectrophotometer (Model 1000, Thermo Scientific). The results were represented as mM TBARS/mg protein.²⁵

Catalase assay (CAT)

For the test, 102 μl of H_2O_2 30% and 100 μl of Triton X-100 were pipetted with 10 μl of the supernatant from the macerate and read at 240 nm in a semi-automatic spectrophotometer (Genesis 8/Spectronic). One CAT unit is defined as 1 μmol of H_2O_2 consumed per minute. The total content of protein in the sample was measured using the Nanodrop Spectrophotometer (Model 1000, Thermo Scientific). The results were presented as CAT units/mg protein.

Reduced glutathione (GSH)

For the analyses, 250 μl of metaphosphoric acid was added to 50 μl of liver supernatant and centrifuged. Afterward, 650 μl of Na_2HPO_4 and 100 μl of the color reagent (5,5'-Dithiobis-2-Nitrobenzoic Acid) were added to 250 μl of the supernatant. A yellow compound was formed and measured in a semi-automatic spectrophotometer at 412 nm (Genesis 8/Spectronic). The total content of protein in the sample was measured using the Nanodrop Spectrophotometer (Model 1000, Thermo Scientific). The results were represented as O.D. 412 $\mu\text{m/g}$ protein.²⁶

Superoxide dismutase (SOD)

The superoxide dismutase was measured with a commercially available kit (RANSOD, Random Labs, UK), according to the manufacturer's instructions. The total content of protein in the sample was measured using the Nanodrop Spectrophotometer (Model 1000, Thermo Scientific). The results were represented as % inhibition/mg protein.

Quantification of mRNA through real-time PCR

Total cellular RNA was extracted from the liver using the Trizol method (Thermo Scientific) according to the manufacturer's instructions. The RNA was resuspended in 20 μl of nuclease-free water (Ambion*) and converted to complementary DNA (cDNA) (GoScript™ Reverse Transcription System Protocol – Promega) according to the protocol indicated by the manufacturer. The final concentration of cDNA was analyzed using the fluorometric method (Qubit® – ThermoFisher – Scientific) using a commercial kit (Qubit® dsDNA HS Assay – Thermo Scientific).

Gene expression was performed in real-time quantitative PCR (Step One Plus – Applied Biosystems) using 16 ng of cDNA. The liver samples were prepared in duplicate and the relative expression of messenger RNA (mRNA) was calculated using the Delta-Delta Ct method ($\Delta\Delta Ct$) using the male control group as a reference. Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was adopted as the reference endogenous gene. A negative control for each primer was used on each plate to check for possible contamination. Amplification reagent measurements were calculated based on the incorporation of the SYBR® Green fluorescence marker (Applied Biosystems) into the double cDNA ribbon for each amplification reaction.

The primers used for real-time PCR were: GAPDH (forward 5' GGGGAGCCAAAAGGGTCATC 3'; reverse 5' GACGCCTGCTTCACCACCTTCTTG 3'), GR (forward 5' GGAATAGGTGCCAAGGGTCT 3'; reverse 5' GAGCACACCAGGCAGAGTTT 3'), PPAR α (forward 5' TGCAAACCTGGACTTGAACG 3'; reverse 5' TGATGTCACAGAACGGCTTC 3') and PPAR γ (forward 5' TGGAATTAGATGACAGTACTTGG 3'; reverse 5' CTCTGTGACGATCTGCCTGAG 3').

Statistical analysis

The normality of data was tested using the Shapiro–Wilk test. Outliers were excluded from analyses. As a normal distribution was found, the data were expressed using mean and standard error of the mean (SEM). Two-way ANOVA with repeated measures was used for the gestational body weight analysis and a one-way ANOVA was used for the offspring body weight analysis. For the evaluation of the interaction between sex (male \times female) and the experimental groups (CON \times PNS \times EX + PNS), a two-way ANOVA was used. Whenever a significant difference was found ($p < .05$), the LSD *post-hoc* test for multiple comparisons was used. For biochemical analysis and PCR data, biological replicates were CON $n = 8$ ($n = 10$ – 14 offspring); PNS $n = 7$ ($n = 7$ – 9 offspring); and EX + PNS $n = 8$ ($n = 13$ – 15 offspring). In all cases, the level of significance was set at 5%. Data were analyzed using SPSS version 18.0 (SPSS Inc., USA).

Results

Pregestational exercise increases neonatal body weight

No effects of prenatal stress and pregestational exercise on maternal body weight were found during gestation, although dams showed a significant body weight gain during pregnancy, as was expected (Fig. 2a). A one-way ANOVA showed that there were no significant effects for groups in the assessment of body weight on PND1 ($F_{(2,20)} = 3.02$; $p = .071$) and PND10 ($F_{(2,20)} = 3.33$; $p = .056$). Results are shown in Fig. 2b and 2c, respectively.

Pregestational exercise and prenatal stress alter body weight in adult life in a sex-dependent manner

A two-way ANOVA revealed significant main effects for group ($F_{(2,48)} = 17.91$; $p = .000002$) and sex ($F_{(1,48)} = 118.00$; $p = .000001$). *Post-hoc* analyses showed that PNS males had a significantly lower body weight when compared to CON ($p = .009$) and EX + PNS ($p = .000011$) groups. Also, at PND 60, EX + PNS males showed a heavier body weight when compared to the CON group ($p = .029$). On the other hand, prenatal stress did not influence female body weight in adult life. However, exercise before pregnancy induced a body weight increase in females when




Fig. 2. Maternal and offspring body weight. (a) Maternal body weight gain during pregnancy was evaluated during the gestational period. Groups were compared by a two-way ANOVA with repeated measures followed by the LSD *post-hoc* test. Data are shown as mean \pm SEM, CON = 8, PNS = 7 and EX + PNS = 8 animals per group. * $p < 0.05$ denotes significant differences between G8 and G15 compared to G1. The offspring body weight was evaluated at (b) PND1 and (c) PND10. Groups were compared by one-way ANOVA. Data are shown as mean \pm SEM, $n = 7$ – 8 animals per group (exact numbers are within the bars). Adult mice's body weight was evaluated at (d) PND60. Groups were compared by a two-way ANOVA followed by the LSD *post-hoc* test. Data are shown as mean \pm SEM, $n = 8$ – 10 animals per group (exact numbers are within the bars). * $p < 0.05$ denotes significant differences compared to the CON group in the same sex; # $p < 0.05$ denotes significant differences compared to the PNS group; & $p < 0.05$ denotes significant sex differences within the same group. CON, control; PNS, prenatal stress from the second week of pregnancy; EX + PNS, physical exercise 3 weeks before gestation and prenatal stress from the second week of pregnancy.

compared to the CON ($p = .001$) and PNS ($p = .002$) groups. As expected, in the comparisons between the sexes, females showed a lower body weight when compared to the males of their respective groups ($p < .0001$). Results are shown in Fig. 2d.

Prenatal stress and pregestational exercise influence the offspring's liver metabolism in a sex-dependent manner

We evaluated the ability of pregestational exercise in preventing the effects of prenatal stress on liver-related biochemical markers.

Table 2. Hepatic oxidative stress markers

Parameters	CON	PNS	EX + PNS
<i>Males</i>			
TBARS (mM/mg protein)	2.089 ± 0.20	2.066 ± 0.31	1.81 ± 0.17
CAT (U/mg protein)	1930 ± 383.6	2980 ± 977.4	1457 ± 201.4
GSH (O.D. 412 μm/g protein)	0.0037 ± 0.0012	0.0036 ± 0.0010	0.0036 ± 0.0010
SOD (%inhibition/mg protein)	0.0163 ± 0.0012	0.0172 ± 0.0019	0.0189 ± 0.0015
<i>Females</i>			
TBARS (mM/mg protein)	2.086 ± 0.44	1.64 ± 0.12	1.453 ± 0.11
CAT (U/mg protein)	3671 ± 850.3	1194 ± 179.0*	1653 ± 175.4*
GSH (O.D. 412 μm/g protein)	0.002 ± 0.0003	0.002 ± 0.0005	0.003 ± 0.0005
SOD (%inhibition/mg protein)	0.0182 ± 0.0024	0.0190 ± 0.0010	0.0203 ± 0.0007

Groups were compared by two-way ANOVA followed by the LSD *post hoc* test. Data are shown as mean ± SEM, $n = 5-9$ animals per group. TBARS, thiobarbituric acid reactive substances; CAT, catalase; GSH, glutathione; SOD, superoxide dismutase; CON, control; PNS, prenatal stress from the second week of pregnancy; EX + PNS, physical exercise three weeks before gestation and prenatal stress from the second week of pregnancy.

* $p < 0.05$ denotes significant differences compared to the CON group in the same sex.




Fig. 3. Biochemical parameters of liver metabolism. Plasma concentrations of (a) glucose, (b) triglyceride and (c) cholesterol were evaluated at PND 60 in males and females. Groups were compared by a two-way ANOVA followed by the LSD *post-hoc* test. Data are shown as mean ± SEM, $n = 5-11$ animals per group (exact numbers are within the bars). * $p < 0.05$ denotes significant differences compared to the CON group in the same sex; # $p < 0.05$ denotes significant differences compared to the PNS group; # $p < 0.05$ denotes significant sex differences within the same group. CON, control; PNS, prenatal stress from the second week of pregnancy; EX + PNS, physical exercise 3 weeks before gestation and prenatal stress from the second week of pregnancy.

No main effects for group, sex and interaction were found for glucose. For the triglyceride, we detected a significant effect for sex ($F_{(1,46)} = 4.12$; $p = .048$). Cholesterol analysis revealed a significant effect for group ($F_{(2,46)} = 4.75$; $p = .013$) and sex ($F_{(1,46)} = 18.42$; $p = .00009$). Fig. 3c shows no significant differences ($p = .057$) between EX + PNS and CON group for plasma cholesterol in males. However, there was a significant increase in the plasma cholesterol in PNS females when compared to CON ($p = .008$). This increase found in the PNS group was prevented by exercising before pregnancy ($p = .021$). When sex effects were analyzed, EX + PNS females had an increased triglyceride secretion ($p = .021$) when compared to the males of the same group (Fig. 3b). Sex differences have also been demonstrated for cholesterol, as CON ($p = .014$) and EX + PNS ($p = .001$) females showed lower cholesterol levels compared to the males from their respective experimental groups (Fig. 3c).

Pregestational exercise does not prevent the effects of prenatal stress on female liver catalase

We also investigated whether pregestational exercise would be capable of preventing possible effects of prenatal stress on oxidative responses in the liver. The liver enzymatic antioxidant defense was evaluated through the activities of CAT, SOD and GSH, besides the

production of TBARS. No significant main effects for group, sex and interaction were observed for the TBARS, GSH and SOD analysis. A significant interaction between group and sex ($F_{(5,34)} = 3.68$; $p = .036$) was shown for catalase and *post-hoc* analysis revealed, in the females, a significant decrease in both PNS ($p = .013$) and EX + PNS ($p = .039$) groups when compared to the CON (Table 2).

Pregestational exercise alters liver gene expression in a sex-dependent manner

Considering the previous results presented, we evaluated GR, PPAR α and PPAR γ mRNA expression in the liver. A significant interaction between group and sex was demonstrated for both GR ($F_{(5,44)} = 3.56$; $p = .037$) and PPAR α ($F_{(5,36)} = 11.37$; $p = .0001$). As for PPAR γ , a significant main effect for group ($F_{(2,32)} = 4.52$; $p = .019$), sex ($F_{(1,32)} = 6.95$; $p = .013$) and interaction ($F_{(5,32)} = 7.51$; $p = .002$) was found. *Post-hoc* analysis showed no significant differences in GR expression between CON and PNS male groups, although a significant GR mRNA expression increase ($p = .006$) was seen when EX + PNS was compared to the PNS group (Fig. 4a). No significant differences were observed for female GR gene expression (Fig. 4a). There was a significant increase in the expression of PPAR α in the liver of PNS ($p = .002$) and




Fig. 4. Hepatic gene expression. (a) Tissue glucocorticoid receptor, (b) peroxisome proliferator-activated receptors α and (c) γ mRNA were measured in males and females. Groups were compared by a two-way ANOVA followed by the LSD *post-hoc* test. Data is shown as mean \pm SEM, $n = 5$ – 11 animals per group (exact numbers are within the bars). * $p < 0.05$ denotes significant differences compared to the CON group in the same sex; & $p < 0.05$ denotes significant differences compared to the PNS group; # $p < 0.05$ denotes significant sex differences within the same group. GAPDH, glyceraldehyde-3-phosphate dehydrogenase; GR, glucocorticoid receptor; PPAR α , peroxisome proliferator-activated receptor α ; PPAR γ , peroxisome proliferator-activated receptor γ ; CON, control; PNS, prenatal stress from the second week of pregnancy; EX + PNS, physical exercise three weeks before gestation and prenatal stress from the second week of pregnancy.

EX + PNS ($p = .036$) males in comparison to the CON group (Fig. 4b). On the other hand, the PPAR α expression significantly decreased in both the PNS ($p = .012$) and EX + PNS ($p = .001$) females when compared to the CON group (Fig. 4b). *Post-hoc* analysis showed a significant decrease in PPAR γ expression in PNS ($p = .001$) and EX + PNS ($p = .000049$) females compared to the CON group (Fig. 4c). Both EX + PNS and PNS did not change the liver PPAR γ expression in males when compared to the CON group (Fig. 4c). When sex effects were analyzed, a significant decrease in the GR mRNA expression ($p = .011$) in EX + PNS females (Fig. 4a) and the PPAR α mRNA expression in both the PNS ($p = .009$) and EX + PNS ($p = .029$) females (Fig. 4b) was observed when compared to the males from the same group. On the other hand, the CON females showed a higher mRNA expression for PPAR α ($p = .002$) and PPAR γ ($p = .000046$) when compared to the CON males (Fig. 4b and 4c, respectively).

Discussion

An adverse environment *in utero* has the ability to affect fetal development and growth, promoting long-term changes in the offspring's metabolism. However, the long-term effects of physical exercise before the gestational period on prenatally stressed offspring, as well as the involvement of sex differences in these responses, have not yet been elucidated. The present study has shown that exercise before pregnancy was able to prevent some of the prenatal stress effects on the metabolic parameters of the offspring in a sex-dependent manner.

Stressful events during the gestational period may disrupt the regulation of metabolic homeostasis.⁹ Both repeated variable prenatal stress and restraint prenatal stress are well-established protocols used in several studies.^{8,13,27,28} Although differences regarding the time and frequency of the repeated restraint stress exposure exist, several studies have demonstrated that its use during pregnancy leads to impaired spatial learning and memory, anxiety, increased corticosterone levels, among others.^{29,30} Moreover, it is well established that exposure to adverse events during the prenatal period is considered to generate alterations in both body weight and composition.^{31–33} Our data showed no effect of stress and pregestational exercise on the weight gain of mothers during gestation. Similarly, Klein *et al.* demonstrated that swimming before gestation does not affect the body weight of pregnant female rats.³⁴ Although swimming has been used as a stressor in repeated variable stress paradigms,^{35,36} studies have also shown its use as a form of exercise, demonstrating positive neural

adaptations, in addition to decreasing corticosterone levels.^{37,38} There is also evidence that mothers who perform physical activities from the beginning of the gestation period and then decrease the exercise levels closer to birth generate significantly heavier children, probably due to the increase in the body fat percentage.³⁹ Using the same stress model during the last 2 weeks of gestation, Vargas *et al.* found no significant differences in the mice's birth weight, corroborating the present findings.²³ Nevertheless, in adulthood, our study showed that prenatal stress decreased body weight only in males in an effect prevented by maternal exercise. Adult females from exercised mothers also presented greater weight compared to the other groups. It is already known that physical activity influences the maintenance of body weight and obesity in rodents.⁴⁰ In a study using rats, treadmill exercise before and during gestation promoted a decrease in the offspring weight at weaning.⁴¹ In mice, the same period of exercise generated an increase in the percentage of lean mass and a decrease of fat mass only in the male adult offspring.⁴² Some differences found between studies may be related to the species used and the type of exercise employed. Moreover, the effects of gestational exercise may be related to changes in body composition, as observed in a study with rats showing sex-dependent effects.⁴³

Considering the body weight results found, we sought to investigate possible liver regulatory mechanisms, since it is a key organ for the control of several metabolic functions. Our data showed that maternal stress caused sex-dependent effects on cholesterol, which is a regulator of lipid organization. Clinical and experimental studies have already demonstrated the beneficial effects of physical exercise on diseases associated with lipid and hepatic alterations,⁴⁴ just as the treadmill running reduced cholesterol in obese and dyslipidemic mice.^{45,46} Increased plasma cholesterol in PNS females had not yet been reported, although maternal stress has already been implicated in the increased lipid accumulation in the liver of prepubertal mice.⁸ It is possible that the beneficial effect of physical exercise prior to gestation in prenatally stressed mice could be associated with both the cholesterol metabolism and sex-dependent mechanisms. Surprisingly, our data showed that males from the exercise group had increased plasma cholesterol, indicating a possible influence of sex hormones since estrogens are already known to be regulators of lipid metabolism.⁴⁷ Although the relationship between exercise and cholesterol levels is well established,^{48–50} we speculate that the association between the withdrawal of exercise before pregnancy and stress may have led to this negative metabolic outcome.

Regarding oxidative stress, our study showed that prenatal stress promotes changes in the antioxidant capacity, observed by a decreased hepatic CAT only in females. At the same time, pre-gestational exercise was not able to influence this effect. CAT is the enzyme responsible for the decomposition of hydrogen peroxide into water and oxygen and it is important for protecting cells from oxidative stress.⁵¹ To date, there have been no studies evaluating the effects of maternal stress on the activity of CAT in the liver of rodents. However, chronic restraint stress is a classic model used to induce alterations in the activity of antioxidant enzymes. It is already known that early-life stress is able to decrease both antioxidant enzymes CAT and SOD levels and thereby increase the production of brain reactive oxygen species in male and female juvenile rats.⁵² Our data demonstrated an effect of prenatal stress in the regulation of oxidative stress only in females, indicating a possible contribution of sex hormones in the antioxidant system response. The type and intensity of physical exercise are also important factors for the production of oxidative stress. A previous study with male mice demonstrated an efficient regulation of antioxidant enzymes and a reduction of oxidative damage in the liver of rodents which were submitted to treadmill exercise compared to sedentary ones.⁵³ Maternal exercise has also been associated with the reversal of deleterious effects of prenatal stress in mice.⁵⁴ Corroborating our findings, a study on rats using maternal swimming before and during gestation did not prevent oxidative changes in the brain of the offspring with neonatal hypoxia-ischemia.⁵⁵

Glucocorticoids are steroid hormones that act predominantly through GR, which are expressed in the hepatic tissue and are involved in the regulation of several energy control processes, besides being associated with the excessive production of oxidative stress.⁵⁶ In addition, it has already been observed that hepatic GR can participate in the regulation of genes involved with cholesterol.^{57,58} In females, our data showed no differences between the groups in the GR gene expression. Similarly, in a murine prenatal model, repeated social stress has been associated with long-term alterations in liver markers of metabolic activity in males and females, although with no changes on the hepatic GR.⁹ Altered GR expression may suggest that exposure to maternal *in utero* glucocorticoids can program the physiology and promote greater susceptibility to metabolic diseases in a sex-dependent manner. The interaction between glucocorticoids and GR is fundamental for the liver homeostatic control, and the decreased expression of these receptors in the tissue has been associated with disorders such as fatty liver and hypercortisolism.⁵⁹ In contrast to the females, pregestational exercise increased the hepatic GR mRNA in males. It has already been observed in rodents that physical exercise induces the increase of circulating glucocorticoid levels, besides increasing the GR expression in the adipose tissue and then inducing lipolysis.⁶⁰ Although the effects of this type of exercise on the offspring's liver have not been addressed, it is possible that our pregestational exercise protocol has acted as a positive modulator of liver GR expression in a sex-dependent manner. Exercising on a treadmill has also been previously considered to be a stressor, although evidence using a similar protocol has already demonstrated a decreased level of glucocorticoids (corticosterone) in exercised mice.⁶¹

Females showed a decreased expression of PPAR α and γ , whereas males showed an increased expression of PPAR α when subjected to stress during pregnancy. In addition, exercise was not able to prevent these effects in both sexes. A study on rats reported that prenatal stress promotes a trend to decrease hepatic mRNA expression of PPAR α and γ only in males.⁹ Moreover, the

study has shown a decreased peroxisome proliferator-activated receptor coactivator 1a (Pgc1a) and transcriptional coactivator (that interacts with PPAR γ) gene expression in females.⁹ Sex differences seem to be an important variable in determining the effects of early life stress on these markers.^{9,62} PPARs play a key role in the metabolism of carbohydrates, lipids and proteins.⁶³ For example, liver disruption of PPAR γ in mice is associated with increased adiposity and hyperlipidemia,⁶⁴ and the lack of PPAR α contributes to the development of steatosis.⁶⁵ The long-term exercise was already related to the increased liver PPAR α in male rats in association with decreased triglyceride and cholesterol.⁶⁶ It is unclear what leads to the increased PPAR α gene expression in PNS males, but the response in the EX + PNS group may be related to hepatic GR levels, since the interaction between these two receptors has already been observed in an inflammatory model.⁶⁷ Very little data are available on mice, related to PPARs gene expression sex-differences. There are studies showing that PPAR-alfa and estrogen receptors share cofactors, suggesting that there is an important role of sex hormones in this response.⁶⁸ In lymphocytes, higher basal levels in PPAR α mRNA and protein in male mice compared to females was demonstrated.⁶⁹ Similarly, a study on rats showed that the expression of PPAR-alfa mRNA and protein levels are higher in male than in female rat livers.⁷⁰ To the best of our knowledge, there are no studies showing sex differences in the expression of PPAR-gamma in mice.

Our study has some limitations, including the lack of a corticosterone assessment in the mothers, which prevented us from investigating if the exercise protocol used was able to alter corticosterone levels. Although it is well-known that both restraint stress has been associated with increased glucocorticoid levels during pregnancy in rodent models,^{71,72} as well as moderate chronic exercise decreases stress-related hormone levels,^{73,74} to the best of our knowledge, there are no studies that have measured corticosterone levels using a similar protocol. Additionally, in order to avoid the use of stressful stimuli to keep mice running, few animals were removed from the study, as they did not adapt to the treadmill. Aiming to minimize the possibility of exercise acting as a stressful stimuli *per se*, all sessions were visually monitored to identify animals with possible difficulties and/or visual signs of distress. Exercising in the last hour of the light cycle may have also influenced the results, as animals are more active during the nocturnal period. PCR was performed using a single reference gene, which would also be considered as a limitation, although GAPDH is stably expressed between groups. Also, direct analysis of the proteins resulting from the mRNA changes detected is also a limitation of the present study, as in mice, the gene expression represents 40% of the variation in protein expression.⁷⁵ Thus, although we did not find significant differences in mRNA expression, different levels of protein could be observed in the liver.

In conclusion, physical exercise before the gestational period prevents the effects of prenatal stress on adult body weight and cholesterol in a sex-dependent manner. However, despite the beneficial metabolic effects found, it was not possible to observe significant differences in the expression of important genes related to the hepatic regulation of this response, such as GR and PPARs. The sex-dependent effects found for stress and exercise suggests the influence of sex steroid hormones, such as estradiol, progesterone and testosterone, although the precise mechanisms still require further investigations.

Acknowledgements. The authors would like to thank CNPq, CAPES and PUCRS for the concession of scholarships.

Financial Support. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001; and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPQ) – Process Number 401178/2016-1.

Conflicts of Interest. None.

Ethical standards. The authors assert that all procedures contributing to this work complied with the ethical standards of the Ethical Committee of Experimental Animals of the Pontifícia Universidade Católica do Rio Grande do Sul and were approved by the institutional committee from this institution.

References

1. Woods SM, Melville JL, Guo Y, Fan MY, Gavin A. Psychosocial stress during pregnancy. *Am J Obstet Gynecol.* 2010; 202, 61.e1–61.e7.
2. Kapoor A, Dunn E, Kostaki A, Andrews MH, Matthews SG. Fetal programming of hypothalamo-pituitary-adrenal function: prenatal stress and glucocorticoids. *J Physiol.* 2006; 572, 31–44.
3. Harris A, Seckl J. Glucocorticoids, prenatal stress and the programming of disease. *Horm Behav.* 2011; 59, 279–289.
4. Tamashiro KL, Moran TH. Perinatal environment and its influences on metabolic programming of offspring. *Phys Behav.* 2010; 100(5), 560–566.
5. Peckett AJ, Wright DC, Riddell MC. The effects of glucocorticoids on adipose tissue lipid metabolism. *Metab Clin Exp.* 2011; 60, 1500–1510.
6. Geer EB, Islam J, Buettner C. Mechanisms of glucocorticoid-induced insulin resistance: focus on adipose tissue function and lipid metabolism. *Endocrin Metab Clin North Am.* 2014; 43, 75–102.
7. Grygiel-Gorniak B. Peroxisome proliferator-activated receptors and their ligands: nutritional and clinical implications – a review. *Nutr J.* 2014; 13, 17.
8. Maeyama H, Hirasawa T, Tahara Y, *et al.* Maternal restraint stress during pregnancy in mice induces 11beta-HSD1-associated metabolic changes in the livers of the offspring. *J Dev Orig Health Dis.* 2015; 6, 105–114.
9. Brunton PJ, Sullivan KM, Kerrigan D, Russell JA, Seckl JR, Drake AJ. Sex-specific effects of prenatal stress on glucose homeostasis and peripheral metabolism in rats. *J Endocrinol.* 2013; 217, 161–173.
10. Zanuso S, Jimenez A, Pugliese G, Corigliano G, Balducci S. Exercise for the management of type 2 diabetes: a review of the evidence. *Acta Diabetol.* 2010; 47, 15–22.
11. Pi H, Liu M, Xi Y, *et al.* Long-term exercise prevents hepatic steatosis: a novel role of FABP1 in regulation of autophagy-lysosomal machinery. *FASEB J.* 2019; 33(11), 11870–11883.
12. Harris JE, Baer LA, Stanford KI. Maternal exercise improves the metabolic health of adult offspring. *Trends Endocrinol Metab.* 2018; 29(3), 164–177.
13. Luft C, Levides IP, Costa MSD, *et al.* Exercise before pregnancy attenuates the effects of prenatal stress in adult mice in a sex-dependent manner. *Int J Dev Neurosci.* 2020; 80, 86–95.
14. Rezaee Z, Marandi SM, Alaei H, Esfarjani F. The effect of preventive exercise on the neuroprotection in 6-hydroxydopamine-lesioned rat brain. *Appl Physiol Nutr Metab.* 2019; 44, 1267–1275.
15. Pitts BL, Whealin JM, Harpaz-Rotem I, *et al.* BDNF Val66Met polymorphism and posttraumatic stress symptoms in U.S. military veterans: protective effect of physical exercise. *Psychoneuroendocrinology.* 2019; 100, 198–202.
16. Stranahan AM, Lee K, Mattson MP. Central mechanisms of HPA axis regulation by voluntary exercise. *Neuromolecular Med.* 2008; 10, 118–127.
17. Herzig KH, Ahola R, Leppaluoto J, Jokelainen J, Jamsa T, Keinänen-Kiukkaanniemi S. Light physical activity determined by a motion sensor decreases insulin resistance, improves lipid homeostasis and reduces visceral fat in high-risk subjects: PreDiabEx study RCT. *Int J Obes (Lond).* 2014; 38, 1089–1096.
18. Sze Y, Brunton PJ. Sex, stress and steroids. *Eur J Neurosci.* 2019; doi: 10.1111/ejn.14615.
19. Weinstock M. Gender differences in the effects of prenatal stress on brain development and behaviour. *Neurochem Res.* 2007; 32, 1730–1740.
20. Viau V, Meaney MJ. Variations in the hypothalamic-pituitary-adrenal response to stress during the estrous cycle in the rat. *Endocrinology.* 1991; 129, 2503–2511.
21. Mauvais-Jarvis F. Sex differences in metabolic homeostasis, diabetes, and obesity. *Biol Sex Differ.* 2015; 6, 14.
22. Wearick-Silva LE, Marshall P, Viola TW, *et al.* Running during adolescence rescues a maternal separation-induced memory impairment in female mice: potential role of differential exon-specific BDNF expression. *Dev Psychobiol.* 2017; 59, 268–274.
23. Vargas MH, Campos NE, de Souza RG, *et al.* Protective effect of early prenatal stress on the induction of asthma in adult mice: sex-specific differences. *Physiol Behav.* 2016; 165, 358–364.
24. Zagron G, Weinstock M. Maternal adrenal hormone secretion mediates behavioural alterations induced by prenatal stress in male and female rats. *Behav Brain Res.* 2006; 175, 323–328.
25. Zanatta A, Viegas CM, Tonin AM, *et al.* Disturbance of redox homeostasis by ornithine and homocitrulline in rat cerebellum: a possible mechanism of cerebellar dysfunction in HHH syndrome. *Life Sci.* 2013; 93, 161–168.
26. Lima KG, Krause GC, da Silva EFG, *et al.* Octyl gallate reduces ATP levels and Ki67 expression leading HepG2 cells to cell cycle arrest and mitochondria-mediated apoptosis. *Toxicol In Vitro.* 2018; 48, 11–25.
27. Kinnunen AK, Koenig JI, Bilbe G. Repeated variable prenatal stress alters pre- and postsynaptic gene expression in the rat frontal pole. *J Neurochem.* 2003; 86, 736–748.
28. Markham JA, Mullins SE, Koenig JI. Periadolescent maturation of the prefrontal cortex is sex-specific and is disrupted by prenatal stress. *J Comp Neurol.* 2013; 521, 1828–1843.
29. Su Q, Zhang H, Dang S, *et al.* Hippocampal protein kinase c gamma signaling mediates the impairment of spatial learning and memory in prenatally stressed offspring rats. *Neuroscience.* 2019; 414, 186–199.
30. Palacios-Garcia I, Lara-Vasquez A, Montiel JF, *et al.* Prenatal stress down-regulates Reelin expression by methylation of its promoter and induces adult behavioral impairments in rats. *PLoS One.* 2015; 10, e0117680.
31. Pankevich DE, Mueller BR, Brockel B, Bale TL. Prenatal stress programming of offspring feeding behavior and energy balance begins early in pregnancy. *Physiol Behav.* 2009; 98, 94–102.
32. Iturra-Mena AM, Arriagada-Solimano M, Luttecke-Anders A, Dagnino-Subiabre A. Effects of prenatal stress on anxiety- and depressive-like behaviours are sex-specific in prepubertal rats. *J Neuroendocrinol.* 2018; 30, e12609.
33. Mueller BR, Bale TL. Impact of prenatal stress on long term body weight is dependent on timing and maternal sensitivity. *Physiol Behav.* 2006; 88, 605–614.
34. Klein CP, Dos Santos Rodrigues K, Hozer RM, *et al.* Swimming exercise before and during pregnancy: promising preventive approach to impact offspring's health. *Int J Dev Neurosci.* 2018; 71, 83–93.
35. Sickmann HM, Skoven C, Bastlund JF, *et al.* Sleep patterning changes in a prenatal stress model of depression. *J Dev Orig Health Dis.* 2018; 9, 102–111.
36. Sickmann HM, Arentzen TS, Dyrby TB, Plath N, Kristensen MP. Prenatal stress produces sex-specific changes in depression-like behavior in rats: implications for increased vulnerability in females. *J Dev Orig Health Dis.* 2015; 6, 462–474.
37. Liu W, Xu Y, Lu J, Zhang Y, Sheng H, Ni X. Swimming exercise ameliorates depression-like behaviors induced by prenatal exposure to glucocorticoids in rats. *Neurosci Lett.* 2012; 524(2), 119–123.
38. Song MK, Kim EJ, Kim JK, Park HK, Lee SG. Effect of regular swimming exercise to duration-intensity on neurocognitive function in cerebral infarction rat model. *Neurol Res.* 2019; 41, 37–44.
39. Finken MJ, van der Voorn B, Heijboer AC, de Waard M, van Goudoever JB, Rotteveel J. Glucocorticoid programming in very preterm birth. *Horm Res Paediatr.* 2016; 85, 221–231.
40. Mifune H, Tajiri Y, Nishi Y, *et al.* Voluntary exercise contributed to an amelioration of abnormal feeding behavior, locomotor activity and ghrelin production concomitantly with a weight reduction in high fat diet-induced obese rats. *Peptides.* 2015; 71, 49–55.

41. Quiclet C, Dubouchaud H, Berthon P, *et al.* Maternal exercise modifies body composition and energy substrates handling in male offspring fed a high-fat/high-sucrose diet. *J Physiol.* 2017; 595, 7049–7062.
42. Carter LG, Lewis KN, Wilkerson DC, *et al.* Perinatal exercise improves glucose homeostasis in adult offspring. *Am J Physiol Endocrinol Metab.* 2012; 303, E1061–E1068.
43. Rosa BV, Blair HT, Vickers MH, *et al.* Moderate exercise during pregnancy in Wistar rats alters bone and body composition of the adult offspring in a sex-dependent manner. *PLoS One.* 2013; 8, e82378.
44. van der Windt DJ, Sud V, Zhang H, Tsung A, Huang H. The effects of physical exercise on fatty liver disease. *Gene Expr.* 2018; 18, 89–101.
45. Cho J, Kim D, Jang J, Kim J, Kang H. Treadmill running suppresses the vulnerability of dopamine D2 receptor deficiency to obesity and metabolic complications: a pilot study. *J Exerc Nutr Biochem.* 2018; 22, 42–50.
46. Ferreira GS, Pinto PR, Iborra RT, *et al.* Aerobic exercise training selectively changes oxysterol levels and metabolism reducing cholesterol accumulation in the aorta of dyslipidemic mice. *Front Physiol.* 2017; 8, 644.
47. Palmisano BT, Zhu L, Eckel RH, Stafford JM. Sex differences in lipid and lipoprotein metabolism. *Mol Metab.* 2018; 15, 45–55.
48. Zhang XL, Zhao N, Xu B, Chen XH, Li TJ. Treadmill exercise inhibits amyloid-beta generation in the hippocampus of APP/PS1 transgenic mice by reducing cholesterol-mediated lipid raft formation. *Neuroreport.* 2019; 30, 498–503.
49. Tarevnic R, Ornellas F, Mandarim-de-Lacerda CA, Aguila MB. Beneficial effects of maternal swimming during pregnancy on offspring metabolism when the father is obese. *J Dev Orig Health Dis.* 2018; 1–5. doi: [10.1017/S2040174418001046](https://doi.org/10.1017/S2040174418001046)
50. Wen S, Jadhav KS, Williamson DL, Rideout TC. Treadmill exercise training modulates hepatic cholesterol metabolism and circulating PCSK9 concentration in high-fat-fed mice. *J Lipids.* 2013; 2013, 908048.
51. Jones DP, Eklow L, Thor H, Orrenius S. Metabolism of hydrogen peroxide in isolated hepatocytes: relative contributions of catalase and glutathione peroxidase in decomposition of endogenously generated H₂O₂. *Arch Biochem Biophys.* 1981; 210, 505–516.
52. Jia N, Sun Q, Su Q, Dang S, Chen G. Taurine promotes cognitive function in prenatally stressed juvenile rats via activating the Akt-CREB-PGC1alpha pathway. *Redox Biol.* 2016; 10, 179–190.
53. da Silva LA, Pinho CA, Rocha LG, Tuon T, Silveira PC, Pinho RA. Effect of different models of physical exercise on oxidative stress markers in mouse liver. *Appl Physiol Nutr Metab.* 2009; 34, 60–65.
54. Bustamante C, Henriquez R, Medina F, Reinoso C, Vargas R, Pascual R. Maternal exercise during pregnancy ameliorates the postnatal neuronal impairments induced by prenatal restraint stress in mice. *Int J Dev Neurosci.* 2013; 31, 267–273.
55. Marcelino TB, de Lemos Rodrigues PI, Miguel PM, Netto CA, Pereira Silva LO, Matte C. Effect of maternal exercise on biochemical parameters in rats submitted to neonatal hypoxia-ischemia. *Brain Res.* 2015; 1622, 91–101.
56. Sato H, Takahashi T, Sumitani K, Takatsu H, Urano S. Glucocorticoid generates ROS to induce oxidative injury in the hippocampus, leading to impairment of cognitive function of rats. *J Clin Biochem Nutr.* 2010; 47, 224–232.
57. Lemke U, Krones-Herzig A, Berriel Diaz M, *et al.* The glucocorticoid receptor controls hepatic dyslipidemia through Hes1. *Cell Metab.* 2008; 8, 212–223.
58. Zollner G, Trauner M. Nuclear receptors as therapeutic targets in cholestatic liver diseases. *Br J Pharmacol.* 2009; 156, 7–27.
59. John K, Marino JS, Sanchez ER, Hinds TD, Jr. The glucocorticoid receptor: cause of or cure for obesity? *Am J Physiol Endocrinol Metab.* 2016; 310, E249–E257.
60. Campbell JE, Fediuc S, Hawke TJ, Riddell MC. Endurance exercise training increases adipose tissue glucocorticoid exposure: adaptations that facilitate lipolysis. *Metab Clin Exp.* 2009; 58, 651–660.
61. Li H, Liang A, Guan F, Fan R, Chi L, Yang B. Regular treadmill running improves spatial learning and memory performance in young mice through increased hippocampal neurogenesis and decreased stress. *Brain Res.* 2013; 1531, 1–8.
62. Paternain L, de la Garza AL, Batlle MA, Milagro FI, Martinez JA, Campion J. Prenatal stress increases the obesogenic effects of a high-fat-sucrose diet in adult rats in a sex-specific manner. *Stress.* 2013; 16, 220–232.
63. Monsalve FA, Pyarasani RD, Delgado-Lopez F, Moore-Carrasco R. Peroxisome proliferator-activated receptor targets for the treatment of metabolic diseases. *Mediat. Inflamm.* 2013; 2013, 549627.
64. Gavrilova O, Haluzik M, Matsusue K, *et al.* Liver peroxisome proliferator-activated receptor gamma contributes to hepatic steatosis, triglyceride clearance, and regulation of body fat mass. *J Biol Chem.* 2003; 278, 34268–34276.
65. Kersten S, Seydoux J, Peters JM, Gonzalez FJ, Desvergne B, Wahli W. Peroxisome proliferator-activated receptor alpha mediates the adaptive response to fasting. *J Clin Invest.* 1999; 103, 1489–1498.
66. Zhang S, Liu Y, Li Q, *et al.* Exercise improved rat metabolism by raising PPAR-alpha. *Int J Sports Med.* 2011; 32, 568–573.
67. Bougarne N, Paumelle R, Caron S, *et al.* PPARalpha blocks glucocorticoid receptor alpha-mediated transactivation but cooperates with the activated glucocorticoid receptor alpha for transrepression on NF-kappaB. *Proc Natl Acad Sci USA.* 2009; 106, 7397–7402.
68. Rando G, Wahli W. Sex differences in nuclear receptor-regulated liver metabolic pathways. *BBA.* 2011; 1812, 964–973.
69. Park HJ, Park HS, Lee JU, Bothwell AL, Choi JM. Gender-specific differences in PPARgamma regulation of follicular helper T cell responses with estrogen. *Sci Rep.* 2016; 6, 28495.
70. Jalouli M, Carlsson L, Ameen C, *et al.* Sex difference in hepatic peroxisome proliferator-activated receptor alpha expression: influence of pituitary and gonadal hormones. *Endocrinology.* 2003; 144, 101–109.
71. Maghsoudi N, Ghasemi R, Ghaempanah Z, Ardekani AM, Nooshinfar E, Tahzibi A. Effect of chronic restraint stress on HPA axis activity and expression of BDNF and Trkb in the hippocampus of pregnant rats: possible contribution in depression during pregnancy and postpartum period. *Basic Clin Neurosci.* 2014; 5, 131–137.
72. Lim R, Fedulov AV, Kobzik L. Maternal stress during pregnancy increases neonatal allergy susceptibility: role of glucocorticoids. *Am J Physiol Lung Cell Mol Physiol.* 2014; 307, L141–L148.
73. Inoue K, Okamoto M, Shibato J, *et al.* Long-term mild, rather than intense, exercise enhances adult hippocampal neurogenesis and greatly changes the transcriptomic profile of the hippocampus. *PLoS One.* 2015; 10, e0128720.
74. Patki G, Li L, Allam F, *et al.* Moderate treadmill exercise rescues anxiety and depression-like behavior as well as memory impairment in a rat model of posttraumatic stress disorder. *Physiol Behav.* 2014; 130, 47–53.
75. Schwanhausser B, Busse D, Li N, *et al.* Global quantification of mammalian gene expression control. *Nature.* 2011; 473, 337–342.